

# Center for Reconstructive Urology

## MEDICATION TO AVOID BEFORE SURGERY

---

The following drugs contain aspirin or other products that can cause increased bleeding during surgery and should not be taken for 7-10 days prior to surgery. Should you need to take something for relief of minor pain, you may take Tylenol.

Advil	Doan's Regular & Extra Strength	Nabumeton
Aleve	Dolobid	Naprelan
Alka Seltzer	Dristan	Naprosyn
Alcohol	Durandyne Tablets	Naproxen
Anacin	Easprin	Norgesic forte
Anaprox	Ecotrin	Orudis
Ansaid	Empirin	Oruvail
Arthrotec	Enbrel	Oxaprozin
Aspirin	Equagesic Tablets	Pepto Bismal Tabs or Liquid
APC	Etodolac	Percodan
BC Tablets or Powder	Excedrin	Persantine
BC Cold Powder	Feldene	Piroxicam
Brufen	Fiorinal	Plavix
Bufferin	Flurbiprofen Sodium	Quagesic
Cama Arthritis Pain Reliever	4 Way Cold Tablets	Relefen
Cataflam	Goody's Headache Powder/Tabs	Robaxisal
Celebrex	Ibuprofen	Rufen
Cephalgesics	Indomethacin	Sine Aid
Clinoril	Indocin	Sulindac
Cogesprin	Ketoprofen	Trandate
Coricidin	Lodine	Trental
Coumadin (Warfarin)	Meclomen	Trilisate
Darvon	Medipren	Vanquish
Darvon with Aspirin	Meloxicam	Vitamin E
Daypro	Midol 200	Voltaren
Diclofenac	Midol PMS Caplets	Zavtrin
Diflunisal	Mobic	Zoprin
Disalcid Tabs or Caps	Motrin	All Herbal supplements

Before stopping any of these medications be sure to consult the physician who ordered them.

**Be advised this is not an all-inclusive list.** If you are unsure whether your medication is an aspirin-based product or an anti-inflammatory, please ask your doctor, nurse or pharmacist.